

## Product Information: Pedialyte AdvancedCare® Plus Powder Sticks

# Pedialyte AdvancedCare® Plus Powder Sticks

Oral rehydration solution for all ages

## Usage

- 1 Pedialyte® AdvancedCare® Plus is formulated with 33% more electrolytes\* to replenish fluids and electrolytes lost during diarrhea and vomiting to help prevent dehydration. Can be used for maintenance of water and electrolytes following corrective parenteral therapy for diarrhea.

## Features

- 1 #1 doctor and pharmacist recommended brand.
- 1 33% more electrolytes\* for higher electrolyte loss.
- 1 Contains PreActiv® prebiotics in the form of fructooligosaccharides (scFOS) that help stimulate the growth of good bacteria in the large intestine.<sup>1</sup>
- 1 When mixed with 500 mL of water:
  - contains the optimal balance of sodium and sugar recommended by experts<sup>2,3</sup> to help prevent dehydration due to diarrhea and vomiting
  - promotes fluid absorption more effectively than common household beverages.
- 1 2 delicious flavours to encourage compliance.
- 1 Low osmolality.
- 1 Gluten-free.
- 1 Kosher.



## Safety Precautions

- 1 Not for parenteral use.
- 1 For infants under 1 year, consult a physician or qualified health care professional before using Pedialyte® AdvancedCare® Plus as infants and young children are at greater risk of dehydration.

\* Sodium electrolytes per equal volume compared to other Pedialyte® products.

\* Sodium electrolytes per equal volume compared to other Pedialyte® products.

<sup>1</sup> Robertfroid, M. et al. British Journal of Nutrition 2010;104:S1-S63.

<sup>2</sup> World Health Organization. Oral Rehydration Salts: Production of the new ORS. 2006.

<sup>3</sup> Leung A, Prince T. Paediatr Child Health. 2006;11(8):527-31.

## Ingredients

### Powder Berry Frost:

MEDICINAL INGREDIENTS: Dextrose, Short-chain Fructooligosaccharides, Citric Acid, Sodium Chloride, Potassium Citrate, Sodium Citrate. NON-MEDICINAL INGREDIENTS: Natural and Artificial Flavours, Calcium Silicate, Acesulfame-Potassium, Sucralose, and FD&C Blue #1.

## Availability

List Number	Item
6696913	Pedialyte® AdvancedCare® Plus Powder Sticks, Berry Frost; 6 x 17-g sticks; 6/case
6697213	Pedialyte® AdvancedCare® Plus Powder Sticks, Strawberry Freeze; 6 x 17-g sticks; 6/case

For more information, contact your Abbott Nutrition sales representative or visit [www.abbottnutrition.ca](http://www.abbottnutrition.ca)

# Pedialyte AdvancedCare® Plus Powder Sticks

Oral rehydration solution for all ages

## Nutrition Information - Powder Berry Frost

	100 mL (at standard dilution)	17 g packet (at standard dilution)
	Value	Value
ENERGY, Cal (kJ)	9.8 (41)	50 (209)
MEDICINAL INGREDIENT:		
Citrate, g (mmol)	0.5 (2.5)	2.57 (13)
Sodium, mg (mmol)	127 (5.9)	650 (30)
Chloride, mg (mmol)	164 (4.9)	840 (25)
Potassium, mg (mmol)	72 (2.0)	370 (10)
Short-chain fructooligosaccharides (scFOS), g	0.14	0.71
Dextrose, g	2.11	

## Product Characteristics

	Value
AT STANDARD DILUTION	
Caloric Density (Cal/mL)	0.1
Protein (% of total energy)	0
Protein Source	N/A
Fat (% of total energy)	0
MCT oil (% of total fats)	N/A
Omega-6:Omega-3 Ratio	N/A
Fat Source	N/A
Carbohydrate (% of total energy)	100
Carbohydrate Source	N/A
Total Cal:g Nitrogen Ratio	N/A
Nonprotein Cal:g Nitrogen Ratio	N/A
Water (g/L)	960
Osmolality (mOsm/kg water)	268
Renal Solute Load (mOsm/L)	130
Recommended Tube Size for Gravity/Pump Feeding (Fr)	N/A

For more information, contact your Abbott Nutrition sales representative or visit [www.abbottnutrition.ca](http://www.abbottnutrition.ca)

# Pedialyte AdvancedCare® Plus Powder Sticks

Oral rehydration solution for all ages

## Preparation

**Directions for Use:**

- 1 Mix the entire contents of 1 stick with 500 mL of drinking water.
- 1 Stir/Shake until completely dissolved. If drinking water is not available, use boiled or bottled water to prepare the solution. Do not boil the solution after it is prepared.
- 1 Tastes best served cold.

**Recommended dose\***

Infants less than 1 year

- 1 Consult a physician before using Pedialyte® AdvancedCare® Plus.

Children 1 year of age

- 1 Give 50-100 mL per episode of diarrhea and/or vomiting, up to a maximum of 1 L over a period of 24 hours.

Children 2–9 years

- 1 Give 100-200 mL per episode of diarrhea and/or vomiting, up to a maximum of 1 L over a period of 8 to 24 hours.

Adults, adolescents and children 10 or more years

- 1 Take 200-400 mL per episode of diarrhea and/or vomiting, or as much as wanted.

\*Attempt to give/take solution as soon as diarrhea begins. Take/give the solution frequently and in small amounts (e.g., 5 mL) gradually increased, as tolerated. If more solution is requested/desired, give/take more, as tolerated.

## Storage & Handling

Store powder sticks at room temperature, avoid extreme temperatures. Refrigerate unused reconstituted product in a sealed container and use within 96 hours or discard.

For more information, contact your Abbott Nutrition sales representative or visit [www.abbottnutrition.ca](http://www.abbottnutrition.ca)