

## Product Information: Similac Pro-Advance® Step 1

For more information, contact your Abbott Nutrition sales representative or visit  
[www.abbottnutrition.ca](http://www.abbottnutrition.ca)

# Similac Pro-Advance® Step 1

with 2'-FL oligosaccharide / DHA- and ARA-fortified

## Usage

- 1 For use as a breast milk substitute in infants who are not exclusively breastfed.
- 1 For oral or tube feeding use.

## Features

- 1 2'-FL\* is structurally identical to 2'-FL human milk oligosaccharide (HMO).
- 1 2'-FL\* is designed to help support infant's developing immune system.
- 1 DHA, lutein and natural vitamin E.‡
- 1 Non-GMO.§
- 1 Easy to digest.¶
- 1 Gluten-free.
- 1 Kosher.
- 1 Halal.



## Safety Precautions

- 1 Not for parenteral use.

\* Not derived from human milk.

‡ RRR-alpha-tocopherol, the most bioactive form of vitamin E.

§ Ingredients not genetically engineered.

¶ Similar to other infant formulas.

For more information, contact your Abbott Nutrition sales representative or visit [www.abbottnutrition.ca](http://www.abbottnutrition.ca)

Abbott Nutrition Abbott Laboratories Co.  
Saint-Laurent, Quebec, Canada H4S 1Z6  
1-866-767-7411

# Similac Pro-Advance® Step 1

with 2'-FL oligosaccharide / DHA- and ARA-fortified

## Ingredients

### Powder Unflavored:

Lactose, Dry or Evaporated Skim Milk, High Oleic Safflower Oil, Whey Protein Concentrate, Soy Oil, Coconut Oil, Fructooligosaccharides, Whey Protein Hydrolysate, M. Alpina Oil (A Source Of ARA), 2'-Fucosyllactose (2'-FL), Schizochytrium Sp. Oil (A Source of DHA), Soy Lecithin, L-Tryptophan, Taurine, MInositol, Cytidine 5'-Monophosphate, Ascorbyl Palmitate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate, Adenosine 5'-Monophosphate, Mixed Tocopherols, L-Carnitine, Lutein, MINERALS (Calcium Carbonate, Potassium Citrate, Potassium Hydroxide, Magnesium Chloride, Sodium Chloride, Ferrous Sulphate, Tricalcium Phosphate, Zinc Sulphate, Copper Sulphate, Manganese Sulphate, Potassium Iodide, Sodium Selenite), VITAMINS (Ascorbic Acid, Choline Bitartrate, D-Alpha-Tocopheryl Acetate, Niacinamide, Calcium D-Pantothenate, Vitamin A Palmitate, Thiamine Chloride Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Beta-Carotene, Phylloquinone, Vitamin D3, D-Biotin, and Cyanocobalamin).

**Allergens:** Contains: Milk, Soy.

## Availability

List Number	Item
6818213	Similac Pro-Advance® Step 1; 859 g can; 6/case

For more information, contact your Abbott Nutrition sales representative or visit [www.abbottnutrition.ca](http://www.abbottnutrition.ca)

Abbott Nutrition Abbott Laboratories Co.  
Saint-Laurent, Quebec, Canada H4S 1Z6  
1-866-767-7411



# Similac Pro-Advance® Step 1

with 2'-FL oligosaccharide / DHA- and ARA-fortified

## Nutrition Information - Powder Unflavored

	100 g (Powder)	100 mL (at standard dilution)	100 Cal
	Value	Value	Value
ENERGY, Cal (kJ)	516 (2160)	68 (280)	100 (418)
Protein, g	11	1.4	2.1
Taurine, mg	35	4.5	6.7
Carnitine, mg	7.6	0.99	1.5
Tryptophan, mg	183	24	35
Fat, g	29	3.7	5.5
Linoleic Acid, g	4	0.53	0.78
Linolenic Acid, g	0.38	0.05	0.075
Arachidonic Acid (ARA), g	0.11	0.014	0.021
Docosahexaenoic Acid (DHA), g	0.055	0.0072	0.011
Carbohydrate, g	56	7.3	10.8
Fibre, g	1.4	0.18	0.27
2'-fucosyllactose (2'-FL), mg	153	20	30
Inositol, mg	30	4	5.9
Lutein, mg	0.058	0.0076	0.011
Nucleotides Equivalents, mg	55	7.2	10.7
Vitamin A, IU	1454	190	282
Vitamin D, IU	309	40	60
Vitamin E, IU	21	2.7	4.0
Vitamin K, mg	0.069	0.009	0.0134
Vitamin C, mg	77	10	15
Thiamine, mg	0.61	0.08	0.12
Riboflavin, mg	0.81	0.11	0.16
Niacin, mg	5.5	0.72	1.06
Vitamin B6, mg	0.31	0.041	0.06
Folic Acid, mg	0.053	0.0069	0.01
Vitamin B12, mg	0.0013	0.00017	0.00026
Pantothenic Acid, mg	2.3	0.3	0.5
Biotin, mg	0.019	0.0025	0.0037
Choline, mg	77	10	15
Sodium, mg	137	18	27
Potassium, mg	609	80	118
Chloride, mg	338	44	65
Calcium, mg	406	53	79
Phosphorus, mg	219	29	42
Magnesium, mg	40	5.2	7.7
Iron, mg	9.2	1.2	1.8
Zinc, mg	3.9	0.5	0.75

For more information, contact your Abbott Nutrition sales representative or visit [www.abbottnutrition.ca](http://www.abbottnutrition.ca)

Abbott Nutrition Abbott Laboratories Co.  
Saint-Laurent, Quebec, Canada H4S 1Z6  
1-866-767-7411



# Similac Pro-Advance® Step 1

with 2'-FL oligosaccharide / DHA- and ARA-fortified

## Nutrition Information - Powder Unflavored

	100 g (Powder)	100 mL (at standard dilution)	100 Cal
	Value	Value	Value
Iodine, mg	0.10	0.013	0.02
Copper, mg	0.41	0.053	0.079
Manganese, mg	0.037	0.0048	0.0071
Selenium, mg	0.010	0.0013	0.002

## Product Characteristics

	Value
AT STANDARD DILUTION	
Caloric Density (Cal/mL)	0.68
Protein (% of total energy)	8.2
Protein Source	Dry or evaporated skim milk, whey protein concentrate, whey protein hydrolysate
Fat (% of total energy)	48.8
Omega-6:Omega-3 Ratio	9.7:1
MCT oil (% of total fats)	0
Fat Source	High oleic safflower oil, soy oil, coconut oil, M. alpina oil, Schizochytrium sp. oil, soy lecithin
Carbohydrate (% of total energy)	43.0
Carbohydrate Source	Lactose, fructooligosaccharides, 2'-fucosyllactose
Total Cal:g Nitrogen Ratio	309:1
Nonprotein Cal:g Nitrogen Ratio	284:1
Osmolality (mOsm/kg water)	303
Renal Solute Load (mOsm/L)	97
Recommended Tube Size for Gravity/Pump Feeding (Fr)	N/A

For more information, contact your Abbott Nutrition sales representative or visit [www.abbottnutrition.ca](http://www.abbottnutrition.ca)

Abbott Nutrition Abbott Laboratories Co.  
Saint-Laurent, Quebec, Canada H4S 1Z6  
1-866-767-7411



# Similac Pro-Advance® Step 1

with 2'-FL oligosaccharide / DHA- and ARA-fortified

## Preparation

### Directions for Preparation and Use:

Your baby's health depends on carefully following these directions. Proper hygiene, preparation, mixing, and storage are important when using infant formula. Failure to follow these directions could result in severe harm. Use as directed by your health care professional.

1. Wash hands thoroughly with soap and water.
2. Sterilize equipment: Place washed bottles, nipples, rings, caps, and all preparation equipment in boiling water and keep at a rolling boil for 2 minutes. Sterilize water for formula: In another pot, boil water at a rolling boil for 2 minutes. Cool to room temperature prior to mixing.
3. Prepare formula using enclosed scoop - see Mixing Guide. Add 1 unpacked level scoop of powder to each 60 mL of cooled, sterilized water. Return dry scoop to container.
4. Cap bottle; shake well; attach nipple.

Once feeding begins, use within 1 hour or discard.

### Mixing Guide

Cooled sterilized water	Unpacked Level Scoop(s)
60 mL	1
120 mL	2
180 mL	3
240 mL	4

## Storage & Handling

Once mixed, refrigerate bottles immediately; use within 24 hours. Store container under normal room temperature conditions. Once opened, store in a cool, dry place with lid tightly closed. Avoid extreme temperatures. Use contents within 1 month. Do not reuse container.

**WARNING:** Powdered infant formulas are not sterile and should not be fed to preterm, low-birth-weight or immunocompromised infants unless directed and supervised by a health care professional. Never use a microwave oven to warm formula; serious burns can result. Test temperature before feeding to avoid scalding baby's mouth. Not for infants/children with galactosemia.

For more information, contact your Abbott Nutrition sales representative or visit [www.abbottnutrition.ca](http://www.abbottnutrition.ca)

Abbott Nutrition Abbott Laboratories Co.  
Saint-Laurent, Quebec, Canada H4S 1Z6  
1-866-767-7411

