

Product Information: Similac Pro-Advance® Step 2

For more information, contact your Abbott Nutrition sales representative or visit
www.abbottnutrition.ca

Similac Pro-Advance® Step 2

with 2'-FL oligosaccharide / DHA- and ARA-fortified

Usage

- 1 For infants and toddlers 6 to 24 months of age.
- 1 Transition formula for infants who are 6 months or older.
- 1 For oral or tube feeding use.

Features

- 1 Specially formulated to help bridge the nutritional gaps that can be associated with the transition to table foods.
- 1 Calcium enriched (50% more calcium than Similac Pro-Advance® Step 1) to support growing infants.
- 1 2'-FL* is structurally identical to 2'-FL found in the breast milk.
- 1 2'-FL* is designed to help support infant's developing immune system.
- 1 DHA, lutein and natural vitamin E.‡
- 1 Non-GMO.§
- 1 Easy to digest.¶
- 1 Gluten-free.
- 1 Kosher.
- 1 Halal.



Safety Precautions

- 1 Not for parenteral use.

* Not derived from human milk.

‡ RRR-alpha-tocopherol, the most bioactive form of vitamin E.

§ Ingredients not genetically engineered.

¶ Similar to other infant formulas.

For more information, contact your Abbott Nutrition sales representative or visit www.abbottnutrition.ca

Abbott Nutrition Abbott Laboratories Co.
Saint-Laurent, Quebec, Canada H4S 1Z6
1-866-767-7411

Similac Pro-Advance® Step 2

with 2'-FL oligosaccharide / DHA- and ARA-fortified

Ingredients

Powder Unflavored:

Lactose, Dry or Evaporated Skim Milk, High Oleic Safflower Oil, Whey Protein Concentrate, Soy Oil, Coconut Oil, Fructooligosaccharides, Whey Protein Hydrolysate, M. Alpina Oil (A Source Of ARA), 2'-Fucosyllactose (2'-FL), Schizochytrium Sp. Oil (A Source of DHA), Soy Lecithin, L-Tryptophan, Taurine, Inositol, Cytidine 5'-Monophosphate, Ascorbyl Palmitate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate, Adenosine 5'-Monophosphate, Mixed Tocopherols, L-Carnitine, Lutein, MINERALS (Tricalcium Phosphate, Calcium Carbonate, Potassium Citrate, Potassium Hydroxide, Magnesium Chloride, Sodium Chloride, Ferrous Sulphate, Zinc Sulphate, Copper Sulphate, Manganese Sulphate, Potassium Iodide, Sodium Selenite), VITAMINS (Ascorbic Acid, Choline Bitartrate, D-Alpha-Tocopheryl Acetate, Niacinamide, Calcium D-Pantothenate, Vitamin A Palmitate, Thiamine Chloride Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Beta-Carotene, Phylloquinone, Vitamin D3, D-Biotin, and Cyanocobalamin).

Allergens: Contains: Milk, Soy.

Availability

List Number	Item
6818313	Similac Pro-Advance® Step 2; 859 g can; 6/case

For more information, contact your Abbott Nutrition sales representative or visit www.abbottnutrition.ca

Similac Pro-Advance® Step 2

with 2'-FL oligosaccharide / DHA- and ARA-fortified

Nutrition Information - Powder Unflavored

	100 g (Powder)	100 mL (at standard dilution)	
	Value	Value	
ENERGY, Cal	516	68	
Protein, g	11	1.4	
Taurine, mg	35	4.5	
Carnitine, mg	7.6	0.99	
Tryptophan, mg	183	24	
Fat, g	29	3.7	
Linoleic Acid, g	4	0.53	
Linolenic Acid, g	0.38	0.05	
Arachidonic Acid (ARA), g	0.11	0.014	
Docosahexaenoic Acid (DHA), g	0.055	0.0072	
Carbohydrate, g	56	7.3	
Fibre, g	1.4	0.18	
2'-fucosyllactose (2'-FL), mg	153	20	
Inositol, mg	30	4	
Lutein, mg	0.058	0.0076	
Nucleotides Equivalents, mg	55	7.2	
Vitamin A, IU	1454	190	
Vitamin D, IU	309	40	
Vitamin E, IU	21	2.7	
Vitamin K, mg	0.069	0.009	
Vitamin C, mg	77	10	
Thiamine, mg	0.61	0.08	
Riboflavin, mg	0.81	0.11	
Niacin, mg	5.5	0.72	
Vitamin B6, mg	0.31	0.041	
Folic Acid, mg	0.053	0.0069	
Vitamin B12, mg	0.0013	0.00017	
Pantothenic Acid, mg	2.3	0.3	
Biotin, mg	0.019	0.0025	
Choline, mg	77	10	
Sodium, mg	137	18	
Potassium, mg	609	80	
Chloride, mg	338	44	
Calcium, mg	609	80	
Phosphorus, mg	304	40	
Magnesium, mg	40	5.2	
Iron, mg	9.2	1.2	
Zinc, mg	3.9	0.5	

For more information, contact your Abbott Nutrition sales representative or visit www.abbottnutrition.ca

Abbott Nutrition Abbott Laboratories Co.
Saint-Laurent, Quebec, Canada H4S 1Z6
1-866-767-7411



Similac Pro-Advance® Step 2

with 2'-FL oligosaccharide / DHA- and ARA-fortified

Nutrition Information - Powder Unflavored

	100 g (Powder)	100 mL (at standard dilution)	
	Value	Value	
Iodine, mg	0.1	0.013	
Copper, mg	0.41	0.053	
Manganese, mg	0.037	0.0048	
Selenium, mg	0.01	0.0013	

Product Characteristics

	Value
AT STANDARD DILUTION	
Caloric Density (Cal/mL)	0.68
Protein (% of total energy)	8.2
Protein Source	Dry or evaporated skim milk, whey protein concentrate, whey protein hydrolysate
Fat (% of total energy)	48.8
Omega-6:Omega-3 Ratio	9.7:1
MCT oil (% of total fats)	0
Fat Source	High oleic safflower oil, soy oil, coconut oil, M. alpina oil, Schizochytrium sp. oil, soy lecithin
Carbohydrate (% of total energy)	43.0
Carbohydrate Source	Lactose, fructooligosaccharides, 2'-fucosyllactose
Total Cal:g Nitrogen Ratio	309:1
Nonprotein Cal:g Nitrogen Ratio	284:1
Osmolality (mOsm/kg water)	303
Renal Solute Load (mOsm/L)	97
Recommended Tube Size for Gravity/Pump Feeding (Fr)	N/A

For more information, contact your Abbott Nutrition sales representative or visit www.abbottnutrition.ca

Abbott Nutrition Abbott Laboratories Co.
Saint-Laurent, Quebec, Canada H4S 1Z6
1-866-767-7411



Similac Pro-Advance® Step 2

with 2'-FL oligosaccharide / DHA- and ARA-fortified

Preparation

Directions for Preparation and Use:

Your baby's health depends on carefully following these directions. Proper hygiene, preparation, mixing, and storage are important when using infant formula. Failure to follow these directions could result in severe harm. Use as directed by your health care professional.

Directions for Preparation and Use:

1. Wash hands thoroughly with soap and water.
2. Sterilize equipment: Place washed bottles, nipples, rings, caps, and all preparation equipment in boiling water and keep at a rolling boil for 2 minutes. Sterilize water for formula: In another pot, boil water at a rolling boil for 2 minutes. Cool to room temperature prior to mixing.
3. Prepare formula using enclosed scoop - see Mixing Guide. Add 1 unpacked level scoop of powder to each 60 mL of cooled, sterilized water. Return dry scoop to container.
4. Cap bottle; shake well; attach nipple.

Once feeding begins, use within 1 hour or discard.

Mixing Guide

Cooled sterilized water	Unpacked Level Scoop(s)
60 mL	1
120 mL	2
180 mL	3
240 mL	4

STORAGE: Once mixed, refrigerate bottles immediately; use within 24 hours. Store container under normal room temperature conditions. Once opened, store in a cool, dry place with lid tightly closed. Avoid extreme temperatures. Use contents within 1 month. Do not reuse container.

WARNING: Powdered infant formulas are not sterile and should not be fed to preterm, low-birth-weight or immunocompromised infants unless directed and supervised by a health care professional. Never use a microwave oven to warm formula; serious burns can result. Test temperature before feeding to avoid scalding baby's mouth.

For more information, contact your Abbott Nutrition sales representative or visit www.abbottnutrition.ca

Abbott Nutrition Abbott Laboratories Co.
Saint-Laurent, Quebec, Canada H4S 1Z6
1-866-767-7411

